


LUPRON DEPOT-PED[®]

(leuprolide acetate for depot suspension)

 Register for Helpful Information About CPP and LUPRON DEPOT-PED



- [Home](#)
- [About CPP](#)
- [About LUPRON DEPOT-PED](#)
- [Starting Therapy](#)
- [Support/Savings](#)
- [FAQs](#)
- [Talking With Your Physician](#)




Does Your Child Have CPP?^{1,2}

Central precocious puberty (CPP) is a condition where girls under 8 years old and boys under 9 begin puberty too early.

- [Signs & symptoms](#)
- [Learn about CPP](#)
- [How is CPP diagnosed?](#)

CPP **OPTIONS** **TREATMENT** **SUPPORT**

LUPRON DEPOT-PED Important Safety Facts

Do Not Take If

- Your child should not receive any formulation of LUPRON DEPOT-PED if he/she has experienced any type of allergic reaction to LUPRON DEPOT-PED or similar drugs.
- Females who are or may become pregnant should not receive any formulation of LUPRON DEPOT-PED.
- LUPRON DEPOT-PED is not for children under 2 years of age.
- View the full Prescribing Information for LUPRON DEPOT-PED.

[Print Safety Facts](#)

Use for LUPRON DEPOT-PED[®] (leuprolide acetate for depot suspension)

LUPRON DEPOT-PED 7.5 mg, 11.25 mg, and 15 mg for 1-month and 11.25 mg and 30 mg for 3-month administration are prescribed for the treatment of children with central precocious puberty (CPP). [More](#)



Ask Your Doctor About 3-Month Dosing²

3-month dosing is designed to offer less-frequent injections—only 4 injections per year for your child vs 12.


[GO >](#)



Get instant savings on LUPRON DEPOT-PED

Your LUPRON DEPOT-PED Instant Savings Card can help you save on every prescription.

[GO >](#)



Talk to a nurse live by phone 1-855-LUPRON-P

Get answers to your questions about LUPRON DEPOT-PED. Learn more.

[GO >](#)

²Based on IMS Health, DDD[™], and NDCHealth[®], NDC Non-Retail[™].

References: 1. Precocious puberty. Mayo Clinic Web site. <http://www.mayoclinic.com/health/precocious-puberty/DS00883>. Accessed March 18, 2014. 2. LUPRON DEPOT-PED [package insert]. 3. Data on file, AbbVie Inc.


You are encouraged to report negative side effects of prescription drugs to the FDA. Visit www.fda.gov/medwatch or call 1-800-FDA-1088.

If you cannot afford your medication, contact www.pparx.org or call the toll-free number 1-888-477-2669.

- [Important Safety Information](#)
- [Glossary of Terms](#)
- [Site Map](#)

- [Prescribing Information](#)
- [Contact Us](#)
- [Privacy Policy](#)
- [Terms of Use](#)

All photos are of models and not persons with central precocious puberty.



Use and Important Safety Information¹

Use for LUPRON DEPOT-PED® (leuprolide acetate for depot suspension)

LUPRON DEPOT-PED 7.5 mg, 11.25 mg, and 15 mg for 1-month and 11.25 mg and 30 mg for 3-month administration are prescribed for the treatment of children with central precocious puberty (CPP).

Doctors may diagnose children with CPP when signs of sexual maturity begin to develop in girls under the age of 8 or boys under the age of 9. Your doctor should perform tests to rule out possible causes of early puberty that would require different treatment (e.g., tumors).

Important Safety Information for LUPRON DEPOT-PED

Your child should not receive any formulation of LUPRON DEPOT-PED if he/she has experienced any type of allergic reaction to LUPRON DEPOT-PED or similar drugs.

Females who are or may become pregnant should not receive any formulation of LUPRON DEPOT-PED.

During the first weeks of treatment, signs of puberty, such as vaginal bleeding, may occur. This is a common initial effect of the drug. Notify your child's doctor if signs/symptoms of puberty continue beyond the second month of treatment.

Convulsions have been observed in patients taking leuprolide acetate, including patients who have a history of seizures, epilepsy, or brain disorders (related to blood vessels, nerves, or tumors), and in patients who are taking medications that have been associated with convulsions. Convulsions have also been reported in patients without any of these conditions.

Your child's initial response to LUPRON DEPOT-PED should be monitored, as well as their continued response during treatment, as clinically needed. If your child receives a 1-month injection, response is assessed 1–2 months after the initial injection. If your child receives a 3-month injection, response is assessed 2–3 months after the initial injection and at month 6. Height and bone age should be checked every 6–12 months in children using LUPRON DEPOT-PED.

It is important that you keep your child's doctor appointments and follow the prescribed injection schedule. Your child's pubertal development could begin again if injections are missed.

The most common side effects with LUPRON DEPOT-PED and similar drugs are pain; acne; injection site reactions, including pain, swelling, and abscess; rash, including a painful rash with fever, blisters/sores, and facial swelling; vaginitis/vaginal bleeding/vaginal discharge; increased weight; altered mood; general pain; headache; fluctuating emotions; and hot flushes/sweating.

After the injection, some pain and irritation is expected; however, if more severe symptoms occur or if any new, unusual, or worsened symptoms develop, contact your child's doctor.

LUPRON DEPOT-PED is not for children under 2 years of age.

LUPRON DEPOT-PED must be administered under the supervision of a physician.

For more information, talk with your child's healthcare provider.

Please see the full [Prescribing Information](#) for LUPRON DEPOT-PED.

Reference: 1. LUPRON DEPOT-PED [package insert].

[Back to the Top](#)

LUPRON DEPOT-PED Important Safety Facts

Do Not Take If

Before Starting

What to Expect

Side Effects

Helpful Resources

[Print Safety Facts](#)

- Your child should not receive any formulation of LUPRON DEPOT-PED if he/she has experienced any type of allergic reaction to LUPRON DEPOT-PED or similar drugs.
- Females who are or may become pregnant should not receive any formulation of LUPRON DEPOT-PED.
- LUPRON DEPOT-PED is not for children under 2 years of age.
- View the full [Prescribing Information for LUPRON DEPOT-PED](#).

Use for LUPRON DEPOT-PED® (leuprolide acetate for depot suspension)

LUPRON DEPOT-PED 7.5 mg, 11.25 mg, and 15 mg for 1-month and 11.25 mg and 30 mg for 3-month administration are prescribed for the treatment of children with central precocious puberty (CPP).

Doctors may diagnose children with CPP when signs of sexual maturity begin to develop in girls under the age of 8 or boys under the age of 9. Your doctor should perform tests to rule out possible causes of early puberty that would require different treatment (e.g., tumors).

LUPRON DEPOT-PED must be administered under the supervision of a physician.

You are encouraged to report negative side effects of prescription drugs to the FDA. Visit www.fda.gov/medwatch or call 1-800-FDA-1088.

If you cannot afford your medication, contact www.pparx.org or call the toll-free number 1-888-477-2669.

[Important Safety Information](#)

[Glossary of Terms](#)

[Site Map](#)

[Prescribing Information](#)

[Contact Us](#)

[Privacy Policy](#)

[Terms of Use](#)



Register for Helpful Information
About CPP and LUPRON DEPOT-PED

About Central Precocious Puberty (CPP)

When are early signs of puberty actually a sign of CPP?¹

Growth spurts, breast development, pubic hair, first periods...[puberty](#) is a time of big changes in your child's body. Normally, you have time to prepare for it. But what happens if your child starts showing signs of puberty at an age you think is much too young?

Central precocious puberty (sometimes called "CPP" or "precocious puberty") is a condition where puberty starts too soon in children—usually **in girls under 8 years old and boys under 9 years old**. If you start seeing early signs of puberty in your child, you should talk with your child's physician.

Learn more about CPP:



What is CPP?



Causes



Signs & Symptoms



Diagnosing



Complications



Treating

Reference: 1. Precocious puberty. Mayo Clinic Web site. <http://www.mayoclinic.com/health/precocious-puberty/DS00883>. Accessed March 18, 2014.

LUPRON DEPOT-PED Important Safety Facts

Do Not Take If

Before Starting

What to Expect

Side Effects

Helpful Resources

- Your child should not receive any formulation of LUPRON DEPOT-PED if he/she has experienced any type of allergic reaction to LUPRON DEPOT-PED or similar drugs.
- Females who are or may become pregnant should not receive any formulation of LUPRON DEPOT-PED.
- LUPRON DEPOT-PED is not for children under 2 years of age.
- View the full [Prescribing Information for LUPRON DEPOT-PED](#).

[Print Safety Facts](#)

Use for LUPRON DEPOT-PED[®] (leuprolide acetate for depot suspension)

LUPRON DEPOT-PED 7.5 mg, 11.25 mg, and 15 mg for 1-month and 11.25 mg and 30 mg for 3-month administration are prescribed for the treatment of children with central precocious puberty (CPP).

Doctors may diagnose children with CPP when signs of sexual maturity begin to develop in girls under the age of 8 or boys under the age of 9. Your doctor should perform tests to rule out possible causes of early puberty that would require different treatment (e.g., tumors).

LUPRON DEPOT-PED must be administered under the supervision of a physician.

[Back to the Top ^](#)

You are encouraged to report negative side effects of prescription drugs to the FDA. Visit www.fda.gov/medwatch or call 1-800-FDA-1088.

If you cannot afford your medication, contact www.pparx.org or call the toll-free number 1-888-477-2669.

[Important Safety Information](#)

[Glossary of Terms](#)

[Site Map](#)

[Prescribing Information](#)

[Contact Us](#)

[Privacy Policy](#)

[Terms of Use](#)



Register for Helpful Information
About CPP and LUPRON DEPOT-PED

About Central Precocious Puberty (CPP)

When are early signs of puberty actually a sign of CPP?¹

Growth spurts, breast development, pubic hair, first periods...[puberty](#) is a time of big changes in your child's body. Normally, you have time to prepare for it. But what happens if your child starts showing signs of puberty at an age you think is much too young?

Central precocious puberty (sometimes called "CPP" or "precocious puberty") is a condition where puberty starts too soon in children—usually **in girls under 8 years old and boys under 9 years old**. If you start seeing early signs of puberty in your child, you should talk with your child's physician.

Learn more about CPP:



[What is CPP?](#)



[Causes](#)



[Signs & Symptoms](#)



[Diagnosing](#)



[Complications](#)



[Treating](#)

Reference: 1. Precocious puberty. Mayo Clinic Web site. <http://www.mayoclinic.com/health/precocious-puberty/DS00883>. Accessed March 18, 2014.

LUPRON DEPOT-PED Important Safety Facts

Do Not Take If

Before Starting

What to Expect

Side Effects

Helpful Resources

- Your child should not receive any formulation of LUPRON DEPOT-PED if he/she has experienced any type of allergic reaction to LUPRON DEPOT-PED or similar drugs.
- Females who are or may become pregnant should not receive any formulation of LUPRON DEPOT-PED.
- LUPRON DEPOT-PED is not for children under 2 years of age.
- View the full [Prescribing Information for LUPRON DEPOT-PED](#).

[Print Safety Facts](#)

Use for LUPRON DEPOT-PED® (leuprolide acetate for depot suspension)

LUPRON DEPOT-PED 7.5 mg, 11.25 mg, and 15 mg for 1-month and 11.25 mg and 30 mg for 3-month administration are prescribed for the treatment of children with central precocious puberty (CPP).

Doctors may diagnose children with CPP when signs of sexual maturity begin to develop in girls under the age of 8 or boys under the age of 9. Your doctor should perform tests to rule out possible causes of early puberty that would require different treatment (e.g., tumors).

LUPRON DEPOT-PED must be administered under the supervision of a physician.

[Back to the Top ^](#)

You are encouraged to report negative side effects of prescription drugs to the FDA. Visit www.fda.gov/medwatch or call 1-800-FDA-1088.

If you cannot afford your medication, contact www.pparx.org or call the toll-free number 1-888-477-2669.

[Important Safety Information](#)

[Glossary of Terms](#)

[Site Map](#)

[Prescribing Information](#)

[Contact Us](#)

[Privacy Policy](#)

[Terms of Use](#)



Register for Helpful Information
About CPP and LUPRON DEPOT-PED

About Central Precocious Puberty (CPP)

When are early signs of puberty actually a sign of CPP?¹

Growth spurts, breast development, pubic hair, first periods...[puberty](#) is a time of big changes in your child's body. Normally, you have time to prepare for it. But what happens if your child starts showing signs of puberty at an age you think is much too young?

Central precocious puberty (sometimes called "CPP" or "precocious puberty") is a condition where puberty starts too soon in children—usually **in girls under 8 years old and boys under 9 years old**. If you start seeing early signs of puberty in your child, you should talk with your child's physician.

Learn more about CPP:



What is CPP?



Causes



Signs & Symptoms



Diagnosing



Complications



Treating

Reference: 1. Precocious puberty. Mayo Clinic Web site. <http://www.mayoclinic.com/health/precocious-puberty/DS00883>. Accessed March 18, 2014.

LUPRON DEPOT-PED Important Safety Facts

Do Not Take If

Before Starting

What to Expect

Side Effects

Helpful Resources

- Your child should not receive any formulation of LUPRON DEPOT-PED if he/she has experienced any type of allergic reaction to LUPRON DEPOT-PED or similar drugs.
- Females who are or may become pregnant should not receive any formulation of LUPRON DEPOT-PED.
- LUPRON DEPOT-PED is not for children under 2 years of age.
- View the full [Prescribing Information for LUPRON DEPOT-PED](#).

[Print Safety Facts](#)

Use for LUPRON DEPOT-PED® (leuprolide acetate for depot suspension)

LUPRON DEPOT-PED 7.5 mg, 11.25 mg, and 15 mg for 1-month and 11.25 mg and 30 mg for 3-month administration are prescribed for the treatment of children with central precocious puberty (CPP).

Doctors may diagnose children with CPP when signs of sexual maturity begin to develop in girls under the age of 8 or boys under the age of 9. Your doctor should perform tests to rule out possible causes of early puberty that would require different treatment (e.g., tumors).

LUPRON DEPOT-PED must be administered under the supervision of a physician.

[Back to the Top ^](#)

You are encouraged to report negative side effects of prescription drugs to the FDA. Visit www.fda.gov/medwatch or call 1-800-FDA-1088.

If you cannot afford your medication, contact www.pparx.org or call the toll-free number 1-888-477-2669.

[Important Safety Information](#)

[Glossary of Terms](#)

[Site Map](#)

[Prescribing Information](#)

[Contact Us](#)

[Privacy Policy](#)

[Terms of Use](#)